

## Western Arctic Caribou Herd Bibliography

### Climate

**Joly, K., P. A. Duffy, and T. S. Rupp. 2012. Simulating the effects of climate change on fire regimes in Arctic biomes: implications for caribou and moose habitat. *Ecosphere* 3 (5): 1-18. Article 36.**

Summary: The amount of area burned in northwest Alaska was modeled using different climate warming scenarios. It is forecasted that fire will increase in the region, which will reduce the amount of high quality habitat by up to 30% within the herd's core winter range. Conversely, moose habitat is forecasted to increase by 20-65%.

**Joly, K., R. R. Jandt, C. R. Meyers, and M. J. Cole. 2007. Changes in vegetative cover on Western Arctic Herd winter range from 1981-2005: potential effects of grazing and climate change. *Rangifer Special Issue 17*: 199-207.**

Summary: Analysis of Western Arctic Caribou Herd winter habitat between 1981 and 2005 revealed a decline in lichens and an increase in grasses overall. These changes are likely to be the result of caribou grazing, fire and possibly climate change.

**Joly, K., D. R. Klein, et al. (2011). "Linkages between large-scale climate patterns and the dynamics of Arctic caribou populations." *Ecography* 34(2): 345-352.**

Summary: Comparison of caribou herd populations with large-scale climate patterns suggest that the Western Arctic Caribou Herd may be showing positive response to the Pacific Decadal Oscillation, which has brought warmer temperatures and wetter conditions in western Alaska and decreased precipitation in the Arctic.

**Joly, K. and D. R. Klein. 2011. Complexity of caribou population dynamics in a changing climate. *Alaska Park Science* 10 (1): 26-31.**

Summary: Caribou have survived previous changes in climate, although may have experienced some localized extinctions. While some effects of climate change may benefit caribou, large changes could be detrimental. This might include changes in habitat that attract other wildlife and more predators. We can expect that caribou populations will change in the future.

### Management

**Callaway, D. (2005). ANILCA and the Western Arctic Caribou Herd Cooperative Management Plan. In: *Alaska Park Science*, December 2005, Vol. 4, No. 2, pp. 22-26.**

Summary: Those who wrote ANILCA appeared to have an awareness of the necessity to integrate local knowledge, values, and cooperation in the framing of a wildlife management regime. When the ADF&G believed the Western Arctic Caribou Herd had dropped to 75,000 caribou in the mid-1970's, local hunters felt these estimates were too low and had missed a lot of animals, so they continued to hunt and exceeded the quota. Employees John Trent and John Cody of ADFG and Dave Spirtes of NPS decided that a new management plan was needed and thus laid the foundation for the Western Arctic Caribou Herd Cooperative Management Plan.

**Dau, J. 2000. Managing reindeer and wildlife on Alaska's Seward Peninsula. Polar Research 19(1):57-62.**

Summary: When reindeer were introduced to the Seward Peninsula in the late 1890's, the Western Arctic Caribou were largely missing from the landscape. As caribou began to return to this area, they have contributed to reduced number of reindeer in herds as reindeer leave and join the migrating caribou or caribou disrupt the reindeer herds and through competition for food and trampling of habitat. The attraction of predators like wolves and brown bears as the number of caribou has increased has had an impact on the reindeer industry. Cooperative management groups have formed recently to try to come up with local solutions to meet the needs of diverse users, including reindeer herders, subsistence and sport hunters and management agencies.

**Fullman, T.J., Joly, K., Ackerman, A. 2017. Effects of environmental features and sport hunting on caribou migration in northwestern Alaska. Movement Ecology 5:4.**

Summary: Influence of physical features (terrain ruggedness, river width, land cover) and sport hunting (hunter camps and transporter aircraft landing sites) on Western Arctic Caribou Herd migration was analyzed for Noatak National Preserve. Caribou avoided rugged terrain and areas with more river, forest, and tall shrubs while selecting for areas dominated by tussock tundra and dwarf shrubs. Migration of caribou through Noatak does not appear to be inhibited by sport hunting activity, though this does not preclude the possibility of temporary effects altering availability of caribou for individual hunters. Caribou exhibited exploratory movement, following predictions of a random walk model. This behavior may facilitate utilization of high-quality forage prior to the onset of winter, especially during mild autumns.

**Klein, D. R., L. Moorehead, et al. (1999). "Contrasts in use and perceptions of biological data for caribou management." Wildlife Society Bulletin 27(2): 488-498.**

Summary: Attitudes and perceptions of hunters and managers of the Western Arctic Caribou Herd were interviewed. Alaskan managers felt that monitoring caribou population dynamics and ranges was their highest priority. Hunters indicated willingness to accept hunting restrictions to help sustain the herd, and accepted the use of radiocollaring and aerial surveys as acceptable methods.

**Rattenbury, K., K. Kielland, et al. (2009). "A reindeer herder's perspective on caribou, weather and socio-economic change on the Seward Peninsula, Alaska." Polar Research 28(1): 71-88.**

Summary: Herder James Noyakuk helped to delineate some of the key factors affecting herders. Weather is more critical now than it may have been 20 years ago, and a single weather event, like delayed freeze-up, early break-up or storms, can have had dramatic effects on herd access and retention. It is difficult to quantify the weather conditions that most affect daily herding plans (visibility, timing of freeze-up and break-up, and snow conditions), whereas air temperature, wind speed and wind direction affect herding indirectly, and did not correlate with how Noyakuk rated weather conditions on herding trips. Responses to extreme weather events are linked to other environmental (caribou, predators, reindeer and range health), economic (depressed international antler prices, low meat and antler sales from small herds, rising fuel and equipment costs, need for non-herding employment) and social (inability to hire crews, inefficiency/dangers of solo herding) stresses) factors.

**Spaeder J., D. Callaway and D. Johnson. 2003. The Western Arctic herd: barriers and bridges to cooperative management. NPS Technical Report NPS/CCSO/UW-2003-01.**

Summary: This report investigated how a cooperative caribou harvest assessment program might contribute to greater trust among Native hunters and federal and state managers. The report detailed a number of case studies and also described and analyzed how cooperative management approaches might be devised to deal with the four general functions of wildlife resource management—research, allocation, regulation, and enforcement.

## **Ecology**

**Ballard, W. B., L. A. Ayres, et al. (1997). "Ecology of wolves in relation to a migratory caribou herd in northwest Alaska." *Wildlife Monographs* (135): 5-47.**

Summary: Studies of wolves and the Western Arctic Caribou Herd during 1987 to 1992 suggested that caribou comprised approximately 51% of wolf diets. Caribou appeared to be the preferred diet of wolves until caribou densities fell below 200 head per ~600 mi<sup>2</sup>, when wolves would switch to preying more moose. Wolves within the range of the Western Arctic Caribou Herd killed 6-7% of this caribou population annually. Wolf predation levels during this period did not appear to strongly limit caribou population growth.

**Dale, B. W., L. G. Adams, et al. (1994). "Functional response of wolves preying on barren-ground caribou in a multiple prey ecosystem." *Journal of Animal Ecology* 63(3): 644-652.**

Summary: Studies during March 1989, & 1990 and November 1990 provided some understanding on how the number and density of caribou on the landscape was affected by wolf predation. From analysis of wolf kill rates, prey selection and prey availability for four wolf packs that overlap with the Western Arctic Caribou Herd, it did not appear that wolves had a regulatory effect on caribou. This was based on mathematical analysis that showed that wolf:caribou ratios were high at low prey densities and that wolves did not increase predation as caribou density increased. Nevertheless, reduction in both wolves and bears would be expected to result in increases in caribou density.

**Dalerum, F., K. Kunkel, et al. (2009). "Diet of wolverines (*Gulo gulo*) in the western Brooks Range, Alaska." *Polar Research* 28(2): 246-253.**

Summary: The diet of approximately 148 wolverine were determined for the period of 1996 to 2002 through stomach and colon content analysis for animals hunted or trapped by locals within the migratory range of the Western Arctic Caribou Herd. Wolverines mostly ate caribou during the winter, many of which may have been killed by other predators and then scavenged by wolverine. When caribou were scarce, wolverine switched to eating mostly moose and were able to stay at a similar fitness level on either diet.

**Dau, J. 2005. Two caribou mortality events in northwest Alaska: possible causes and management implications. – *Rangifer Special Issue* 16: 37-50.**

Summary: After receiving reports from hunters of small numbers of dead caribou near Pt. Hope during the winters of 1994-1995 and 1999-2000, a search was done to count carcasses and sample them to determine what had happened. It appeared that many caribou were malnourished and a high number of calves had died. There was no clear evidence of toxins, but analysis of weather data suggested that wind may have produced hardened snow that could have made it hard for caribou to feed and very low temperatures further stressed the caribou.

These deaths did not appear to affect the overall population of the herd but had an impact on local groups.

**Dau, J. 2009. Western Arctic Caribou Herd. – In Harper, P. (ed.). Caribou management\_ report\_ of survey –inventory activities, 1 July 2006–30 June 2008. Alaska Department of Fish and Game. Juneau, Alaska, USA**

Summary: Censuses of the WAH were conducted in 2007 and 2009. The July 2007 photos census produced a minimum estimate of 377,000 caribou, but may have been underestimated due to poor lighting and unaccounted for small groups. In 2009, the herd was more completely aggregated and overall good conditions for the census provide confidence in an estimate of approximately 401,000 caribou. During June calving surveys, we observed 65 calves:100 cows in 2006; 73:100 in 2007 and 70:100 in 2008. June, fall and spring calf:cow ratios have declined at similar rates during 1982-2008. Pregnant cows and some nonmaternal caribou begin migrating toward the calving grounds in the Utukok River uplands in April.

**Gerlach, S. C., L. K. Duffy, et al. (2006). "An exploratory study of total mercury levels in archaeological caribou hair from northwest Alaska." *Chemosphere* 65(11): 1909-1914.**

Summary: Examination of caribou hair from an archaeological site near Deering from about 860 years ago suggest similar levels of mercury then as is found now. These historical data will be important to look for changes as we monitor levels of mercury in subsistence foods in the future.

**Hong, G. H., M. Baskaran, et al. (2011). "Anthropogenic and natural radionuclides in caribou and muskoxen in the Western Alaskan Arctic and marine fish in the Aleutian Islands in the first half of 2000s." *Science of the Total Environment* 409(19): 3638-3648.**

Summary: Caribou samples from the Western Arctic Caribou Herd collected between 1998 and 2006 were analyzed for radionuclides from natural and human-activity related sources. Levels found in caribou were similar to those found in the plants they eat, and levels were low and did not appear to pose a health threat to subsistence communities during this time.

**Joly, K., R. R. Jandt, et al. (2009). "Decrease of lichens in Arctic ecosystems: the role of wildfire, caribou, reindeer, competition and climate in north-western Alaska." *Polar Research* 28(3): 433-442.**

Summary: A review of research reveals signs of change in Arctic tundra ecosystems. Factors known to be affecting these changes include wildfire, disturbance by caribou and reindeer, differential growth responses of vascular plants and lichens, and associated competition under climate warming scenarios. These factors suggest a reduction in lichen during recent decades.

**Joly, K., S. K. Wasser, and R. Booth (2015) Non-invasive assessment of the interrelationships of diet, pregnancy rate, group composition, and physiological and nutritional stress of barren-ground caribou in late winter. *PLoS One* 10 (6): e0127586. doi:10.1371/journal.pone.0127586.**

Summary: Fecal samples were collected and analyzed to assess the late winter diets, rates of pregnancy, sex ratios and condition of Western Arctic Herd caribou. Diets were dominated by lichens. Pregnancy rates fell within the expected range for a declining here.

## Winter

**Joly, K. 2011. Modeling influences on winter distribution of caribou in northwestern Alaska through use of satellite telemetry. Rangifer Special Issue 19: 75-85.**

Summary: Satellite collars fitted to caribou revealed that that caribou moved significantly less during mid-winter than early- or late-winter and that cows moved significantly more in April than bulls due to their earlier departure on their spring migration. Cows avoided forested areas, highlighting the importance of tundra habitats, and selected for dwarf shrub, with relatively high lichen cover, and sedge habitat types. Bulls selected for dryas, coniferous forest and dwarf shrub habitats but against lowland sedge, upland shrub and burned tundra.

**Joly, K., P. Bente, et al. (2007). "Response of overwintering caribou to burned habitat in northwest Alaska." Arctic 60(4): 401-410.**

Summary: Satellite collars fitted to caribou revealed that caribou were more likely to select burned areas in the late fall and early spring than midwinter. They appear to particularly avoid burns estimated to be 26-55 years old, likely due to the reduction in lichen in burned areas.

**Joly, K., F. S. Chapin, et al. (2010). "Winter habitat selection by caribou in relation to lichen abundance, wildfires, grazing, and landscape characteristics in northwest Alaska." Ecoscience 17(3): 321-333.**

Summary: We found that lichen abundance was more than 3 times greater at locations used by caribou than found at random. The current winter range does not appear to be overgrazed as a whole, but continued high grazing pressure and consequences of climate change on plant community structure might degrade its condition. Within the current winter range, lichen abundance was more than 4 times greater at unburned locations than at recently (< 58 y) burned locations. The historic winter range has low lichen abundance, likely due to sustained grazing pressure exerted by the herd, which suggests that range deterioration can lead to range shifts.

**Joly, K., M. J. Cole, et al. (2007). "Diets of Overwintering Caribou, Rangifer tarandus, Track Decadal Changes in Arctic Tundra Vegetation." Canadian Field-Naturalist 121(4): 379-383.**

Summary: Winter diets of the Western Arctic Caribou Herd from 1995/1996 and 2005 were determined based on fecal analysis. Diets tracked changes in vegetation observed across 10 years, with decreased lichen and increased sedges and shrubs.

**Joly, K., P. A. Duffy, and T. S. Rupp. 2012. Simulating the effects of climate change on fire regimes in Arctic biomes: implications for caribou and moose habitat. Ecosphere 3 (5): 1-18. Article 36.**

Summary: The amount of area burned in northwest Alaska was modeled using different climate warming scenarios. It is forecasted that fire will increase in the region, which will reduce the amount of high quality habitat by up to 30% within the herd's core winter range. Conversely, moose habitat is forecasted to increase by 20-65%.

**Saperstein, L. B., 1996. Winter forage selection by barren-ground caribou: Effects of fire and snow. Rangifer Special Issue, 9:237-238.**

Summary: Both long- and short-term consequences should be considered when examining the effects of fire on the foraging behavior of caribou. Post-fire increases in protein content, digestibility, and availability of cotton grass, *E. vaginatum*, make burned tussock tundra an

attractive feeding area for caribou in late winter. These benefits are likely short-lived, however. Lowered availability of lichens and increased relative frequency of bryophytes will persist for a much longer period.

**Skoog, R.O., 1968. Ecology of the caribou (*Rangifer tarandus granti*) in Alaska. Ph.D. dissertation. University California, Berkeley. 699 pp., RO. 1968. Ecology of the caribou (*Rangifer tarandus granti*) in Alaska. Ph.D. dissertation. University California, Berkeley. 699pp.**

Summary: Alaska's caribou population consists of six regional sub-populations and eleven herds, typically centered around high quality habitat. As the number of caribou in an area increase, they begin to move into areas of lower quality habitat and some may eventually leave the herd and go into other regions. A historical view of caribou herds shows that many animals moved between regions, especially when there were more than 5 to 10 animals per square mile. It appears that there was enough forage in many regions to support more caribou when this study was completed in the 1960's.

## Resource Development

**Magdanz, James S., Joshua Greenberg, Joseph M. Little, and David S. Koster. 2016. The Persistence of Subsistence: Wild Food Harvests in Rural Alaska, 1983-2013. *Social Science Research Network* 2779464. [http://papers.ssrn.com/sol3/papers.cfm?abstract\\_id=2779464](http://papers.ssrn.com/sol3/papers.cfm?abstract_id=2779464)**

Summary: Explores Alaska's subsistence economy using community-level demographic, economic, and harvest data from a large sample of rural households, evaluating trends over time, identifying factors associated with harvests and personal incomes, and modeling productivity. With time as the only factor, models suggest community populations and wild food harvests are not changing, while real personal incomes are gradually declining, contradicting popular narratives of burgeoning rural indigenous populations, diminishing subsistence and growing cash-dependence. Multiple regression models find evidence of modest harvest declines over time in remote areas; geographic location and accessibility are highly influential. Using propensity score matching, models show that road building is expected to result in significant declines in harvests, but no significant changes in incomes, contradicting state narratives of positive rural development impacts.

**Wilson, R.R., D.D. Gustine, and K. Joly. 2014. Evaluating Potential Effects of an Industrial Road on Winter Habitat of Caribou in North-Central Alaska. *Arctic* 67(4): 472-482.**

Summary: Worldwide, some caribou (*Rangifer tarandus*) populations are experiencing declines due partially to the expansion of industrial development. Caribou can exhibit behavioral avoidance of development, leading to indirect habitat loss, even if the actual footprint is small. Thus, it is important to understand before construction begins how much habitat might be affected by proposed development. In northern Alaska, an industrial road that has been proposed to facilitate mining transects a portion of the Western Arctic caribou herd's winter range. To understand how winter habitat use might be affected by the road, we estimated resource selection patterns during winter for caribou in a study area surrounding the proposed road. We assessed the reductions of habitat value associated with three proposed routes at three distance thresholds for disturbance. High-value winter habitat tended to occur in locally rugged areas that have not burned recently and have a high density of lichen and early dates of spring snowmelt. We found that 1.5% to 8.5% (146–848 km<sup>2</sup>) of existing high-value winter habitat in our study area might be reduced in quality. The three alternative routes were only

marginally different. Our results suggest that the road would have minimal direct effects on high-value winter habitat; however, additional cumulative impacts to caribou (e.g., increased access by recreationists and hunters) should be considered before the full effects of the road can be estimated.

**Wilson, R.R., L.S. Parrett, K. Joly, and J.R. Dau. 2016. Effects of roads on individual caribou movements during migration. *Biological Conservation* 195: 2-8.**

Summary: Long distance migrations by large mammals are increasingly imperiled by human development. We studied autumn migratory patterns of caribou (*Rangifer tarandus*) in relation to an industrial road in northwestern Alaska. We built null movement models to determine the expected time to cross the road if caribou movements were not affected by the road. We then identified individuals that took longer to cross than expected (slow crossers) and those that did not differ from that expected from the null model (normal crossers). We identified eight as slow and 20 as normal crossers. Slow crossers took an average of  $33.3 \pm 17.0$  ( $\pm$ SD) days to cross the road compared to  $3.1 \pm 5.5$  days for normal crossers. Slow crossers had an average crossing date of 8 Nov.  $\pm 7.7$  days versus 25 Oct.  $\pm 20.6$  days for normal crossers. Movement rates of the two classes did not differ before crossing the road, but slow crossers moved N1.5 times as fast as normal crossers after crossing the road. Movement patterns were partially explained by environmental attributes, but were most strongly affected by how far a caribou was from the road and whether it was classified as slow or normal crosser. While avoidance is an important aspect of the effects of roads on populations, our results show the importance of other factors, such as how long individuals are delayed in crossing when assessing the influence of development on wildlife.

**Wolfe, R.J. and R.J. Walker. 1986. *Impacts of roads and settlement entry on subsistence in Alaska*. Proceedings of the Annual Meeting of Alaska Anthropological Association: Fairbanks.**

**Wolfe, R.J. and R.J. Walker. 1987. *Subsistence economies in Alaska: productivity, geography, and development impacts*. *Arctic Anthropology* 24(2), pages 56–81.**

## Subsistence Use

**Andersen, David B., Caroline Brown, Robert Walker, and Gretchen Jennings. 2004. The 2001–2002 Harvest of Moose, Caribou, and Bear in Middle Yukon and Koyukuk River Communities. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 278.**

**Andersen, David B., Charles J. Utermohle, and Louis Brown. 1998. The 1997–1998 Harvest of Moose, Caribou, and Bear in Middle Yukon and Koyukuk River Communities, Alaska. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 245.**

**Andersen, David B., Charles J. Utermohle, and Louis Brown. 2000. The 1998–1999 Harvest of Moose, Caribou, and Bear in Ten Middle Yukon and Koyukuk River Communities. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 251.**

**Andersen, David B., Charles J. Utermohle, and Gretchen Jennings. 2001. The 1999–2000 Harvest of Moose, Caribou, and Bear in Ten Middle Yukon and Koyukuk River Communities. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 262.**

**Braund, Steven R., Karen Brewster, Lisa Moorehead, Timothy P. Holmes, and John A. Kruse. 1993. The North Slope Subsistence Study: Barrow, 1987, 1988, 1989. Anchorage: Technical Report No. 149, OCS Study, MMS 91-0086, submitted to U.S. Department of the Interior Minerals Management Service, for contract no. 14-12-0001-30284. Stephen R. Braund & Associates with University of Alaska Anchorage Institute of Social and Economic Research.**

**Braund, Steven R., Eric Loring, Lisa Moorehead, David C. Burnham, and John A. Kruse. 1993. The North Slope Subsistence Study: Wainwright, 1988, 1989. Submitted to the US Department of Interior, Minerals Management Service, Alaska OCS Region, Anchorage, Alaska. Anchorage: Stephen R. Braund & Associates with Institute of Social and Economic Research, University of Alaska Anchorage.**

**Braem, N.M., S. Pedersen, J. Simon, D.S. Koster, T. Kaleak, P. Leavitt, J. Patkotak, and P. Neakok. 2011. Monitoring of caribou harvests in the National Petroleum Reserve in Alaska: Atqasuk, Barrow, and Nuiqsut, 2003–2007. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 361, Fairbanks.**

**Braem, N.M., 2011. Subsistence wildlife harvests in Deering, Alaska, 2007-2008. Alaska Department of Fish and Game Division of Subsistence, Special Publication No. SP2010-002, Anchorage.**

**Braem, N.M., 2011. Subsistence wildlife harvests in Noorvik, Shungnak and White Mountain, Alaska, 2008-2009. Alaska Department of Fish and Game Division of Subsistence, Special Publication No. SP2011-003, Fairbanks.**

**Braem, N.M., 2012. Subsistence wildlife harvests in Ambler, Buckland, Kiana, Kobuk, Shaktoolik and Shishmaref, Alaska, 2009-2010. Alaska Department of Fish and Game Division of Subsistence, Special Publication No. SP2012-003, Fairbanks.**

**Braem, N.M.; James S. Magdanz; David S. Koster; Patricia Fox. 2013. Subsistence harvests in Northwest Alaska: Selawik, 2010–2011. ADF&G Division of Subsistence, Technical Paper No. 389.**

**Braem, N.M. and M. Kostick. 2014. Subsistence wildlife harvests in Elim, Golovin, Kivalina, Koyuk, Noatak, and Wales, Alaska, 2010-2011. Alaska Department of Fish and Game Division of Subsistence, Special Publication No. SP2012-004, Fairbanks.**

**Braem, N.M., E.H. Mikow, S.J. Wilson, and M.L. Kostick. 2015. Wild Food Harvests in 3 Upper Kobuk River Communities: Ambler, Shungnak, and Kobuk, 2012–2013. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 402, Fairbanks.**

**Braem, N.M., E.H. Mikow, and M.L. Kostick, editors. 2017. Chukchi Sea and Norton Sound Observation Network: Harvest and Use of Wild Resources in 9 Communities in Arctic Alaska, 2012–2014. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 403, Fairbanks.**



**Braem, Nicole M. 2017. Revised Options for Amounts Reasonably Necessary for Subsistence Uses of the Teshekpuk Caribou Herd. Alaska Department of Fish and Game Division of Subsistence, Special Publication No. BOG 2017-02, Fairbanks.**

**Brower, R., and T. Opie. 1996. North Slope Borough Subsistence Harvest Documentation Project: Data for Anaktuvuk Pass, Alaska for the Period July 1, 1994 to June 30, 1995. Barrow: North Slope Borough, Department of Wildlife Management.**

**Brower, R., and T. Opie. 1997. North Slope Borough Subsistence Harvest Documentation Project: Data for Atqasuk, Alaska for the Period July 1, 1994 to June 30, 1995. Barrow: North Slope Borough, Department of Wildlife Management.**

**Brower, R., and T. Opie. 1998. North Slope Borough Subsistence Harvest Documentation Project: Data for Nuiqsut, Alaska for the Period July 1, 1994 to June 30, 1995. Barrow: North Slope Borough, Department of Wildlife Management.**

**Brown, Caroline L., Robert Walker, and Susan B. Vanek. 2004. The 2002-2003 Harvest of Moose, Caribou, and Bear in Middle Yukon and Koyukuk River Communities. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 280.**

**Brown, C.L., N.M. Braem, M.L. Kostick, A. Trainor, L.J. Slayton, D.M. Runfola, E.H. Mikow, H. Ikuta, C.R. McDevitt, J. Park, and J.J. Simon. 2016. Harvests and uses of wild resources in 4 Interior Alaska communities and 3 Arctic Alaska communities. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 426, Fairbanks.**

**Fuller, Allen, S., and John George C. 1997. "Evaluation of Subsistence Harvest Data from the North Slope Borough 1993 Census for Eight North Slope Villages: For the Calendar Year 1992." Barrow: North Slope Borough, Department of Wildlife Management.**

**Galginaitis, Michael C. 1990. "Subsistence Resource Harvest Patterns: Nuiqsut." Anchorage: Submitted to the US Department of Interior, Minerals Management Service, Alaska OCS Region.**

**Galginaitis, Michael C., Claudia Chang, Kathleen M. MacQueen, Albert A. Dekin Jr., and David Zipkin. 1984. "Ethnographic Study and Monitoring Methodology of Contemporary Economic Growth, Sociocultural Change and Community Development in Nuiqsut, Alaska." Anchorage: Social and Economic Studies Program Technical Report No. 96. Submitted to the US Department of Interior, Minerals Management Service, Alaska OCS Region.**

**Georgette, S. 1994. Summary of Western Arctic Caribou herd overlays (1984-92) and comparison with harvest data from other sources. Alaska Department of Fish and Game, Division of Subsistence: Juneau.**

**Georgette, S. 1999. Subsistence harvests in Northwest Alaska: Caribou, Moose, Bear, Wolf, and Wolverine May 1998 through April 1999. Alaska Department of Fish and Game, Division of Subsistence: Kotzebue.**

**Georgette, S. and H. Loon 1988. *The Noatak River: Fall caribou hunting and airplane use.* Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 162.**

**Georgette, S. and H. Loon. 1993. *Subsistence use of fish and wildlife in Kotzebue, a Northwest Alaska regional center.* Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 167: Juneau.**

**Georgette, S., K. Persons, and A. Ahmasuk. 2002. *Subsistence Wildlife Harvests in Four Communities on the Western Seward Peninsula, Alaska 2000-2001.* Alaska Department of Fish and Game Division of Subsistence and Kawerak, Inc.: Kotzebue.**

**Georgette, S., K. Persons, E. Shiedt, and S. Tahbone. 2004. *Subsistence wildlife harvests in five northwest Alaska communities, 2001–2003.* Alaska Department of Fish and Game, Maniilaq Association, and Kawerak, Inc.**

**Georgette, S., A. Ahmasuk, K. Persons, E. Shiedt, and E. Trigg. 2005. *Subsistence wildlife harvests in three northwest Alaska communities, 2003–2004.* Alaska Department of Fish and Game, Maniilaq Association, and Kawerak, Inc.**

**Godduhn, A.R., N.M. Braem, and M.L. Kostick 2014. *Subsistence Wildlife Harvests in Kotzebue, Alaska, 2012–2013.* Alaska Department of Fish and Game Division of Subsistence, Special Publication No. SP2014-03: Fairbanks**

**Guettabi, M., J. Greenberg, J. Little, and K. Joly. 2016. *Evaluating potential economic effects of an industrial road on subsistence in north-central Alaska.* *Arctic* 69 (3): 305-317.** A new road has been proposed to provide access to this region and the Ambler Mining District from north-central Alaska. To evaluate how it might affect subsistence harvest, we used models to identify factors related to subsistence production at the household level. We found substantial differences in these factors between communities near the proposed road and a comparable set of road accessible communities outside the region. Total subsistence production of study area communities was 1.8 to 2.5 times greater than those outside it. If the road was opened to the public and subsistence harvest patterns for study area communities changed to mirror existing communities on the existing road system as a result of the road, the financial cost would be \$6,900 – \$10,500 per household (assuming a \$8/lb “replacement” cost for subsistence harvests). Taken together, our results suggest that the proposed road should be expected to substantially impact subsistence production in communities that are not currently connected to the road system.

**Halas, G, and G. Kofinas. 2015. *Community Report: Caribou Migration, Subsistence Hunting, and User Group Conflicts in Northwest Alaska.* UAF School of Natural Resources & Extension, AFES Miscellaneous Report 2015-06.**

Summary: This community report presents key findings of a research project of the University of Alaska Fairbanks, the National Park Service, and the Native Village of Noatak that studied the links between caribou, interactions of local and non-local hunters, and changes to subsistence caribou hunting. (<http://www.uaf.edu/snre/research/publications/miscellaneous-publication/>)

**Halas, Gabriela. 2015. *Caribou migration, subsistence hunting, and user group conflicts in northwest Alaska: A traditional knowledge perspective.* School of Natural Resources and**

**Extension, University of Alaska Fairbanks.**

Summary: In the last decade there has been an increase in caribou hunting activities by non-local hunters and commercial operators in Noatak's traditional hunting lands, including the Noatak National Preserve.

This thesis acts as the full report from a research project of the University of Alaska Fairbanks, the National Park Service, and the Native Village of Noatak that studied the links between caribou, interactions of local and non-local hunters and transports, and changes to subsistence caribou hunting.

**Heller, C.A. and E.M. Scott 1967. *The Alaska dietary survey, 1956–1961*. U. S. Department of Health: Education, and Welfare Nutrition and Metabolic Disease Section, Arctic Health Research Center: Anchorage**

**Magdanz, J.S., N.S. Braem, B.C. Robbins, and D.S. Koster 2010. *Subsistence harvests in Northwest Alaska, Kivalina and Noatak, 2007*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 354: Kotzebue.**

**Magdanz, J.S., C.J. Utermohle, and R.J. Wolfe 2002. *The production and distribution of wild food in Wales and Deering, Alaska*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 259: Kotzebue.**

**Magdanz, J.S., R.J. Walker, and R.R. Paciorek 2004. *The subsistence harvests of wild foods by residents of Shungnak, Alaska, 2002*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 279: Juneau**

**Marcotte, James R., and Terry L. Haynes. 1985. "Contemporary Resource Use Patterns in the Upper Koyukuk Region, Alaska." Technical paper 93. Fairbanks: Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp093.pdf>.**

**Mikow, E., Nicole M. Braem, Marylyne Kostick. 2014. *Subsistence Wildlife Harvests in Brevig Mission, Deering, Noatak, and Teller, Alaska, 2011–2012*. Alaska Department of Fish and Game Division of Subsistence, Special Publication No. -002.**

**Mikow, E.H. and M.L. Kostick 2016. *Subsistence wildlife harvests in Kotzebue, Alaska, 2013–2014*. Alaska Department of Fish and Game Division of Subsistence, Special Publication No. 2016-2: Fairbanks**

**Patterson, A. 1974. *Subsistence harvests in five native regions: for the Joint Federal-State Land Use Planning Commission for Alaska*. Resource Planning Team: Anchorage.**